



Bern University, CERN and JINR Dubna

HadAtom05

Workshop on Hadronic Atoms,
University of Bern, Switzerland
February 15 and 16, 2005



edited by

L. Afanasyev¹, G. Colangelo² and J. Schacher³

¹ *Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia*

² *Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland*

³ *Laboratory for High-Energy Physics, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland*

Abstract

These are the proceedings of the workshop “HadAtom05”, held at the University of Bern, Switzerland, February 15 and 16, 2005. The main topics of the workshop concerned the physics of hadronic atoms, and in this context recent results from experiments and theory were presented. These proceedings contain the list of participants, the scientific program and a short contribution from each speaker.

1 Introduction

The workshop “HadAtom05” was held at the University of Bern, Switzerland, on February 15 and 16, 2005. This was already the sixth regular workshop of the HadAtom type. This workshop series has originally been inspired by the latest experimental and theoretical progress achieved in the investigation of the bound states of strongly interacting particles, the hadronic atoms. The previous workshops were held in Dubna (1998), Bern (1999, 2001), at CERN (2002) and in Trento (2003). The present workshop covered the following topics:

- Hadronic atoms, in particular their
 - Production
 - Interaction with matter
 - Energy levels
 - Decays
- Meson-meson and meson-baryon scattering
- Lattice calculations and low energy effective theory of QCD
- Chiral Perturbation Theory and nuclear many-body systems
 - Nuclear Matter
 - Pionic deuterium and pionic hydrogen
- Experiments
 - DIRAC at CERN
 - DEAR/SIDDHARTA at DAFNE
 - Pionic Hydrogen Collaboration at PSI
 - Hadronic atoms at J-PARC and GSI
 - K decays with NA48 at CERN

About 40 physicists took part in the workshop, and 23 talks were presented. As for the previous workshops [1,2,3,4,5], we publish a collection of abstracts of the presentations, containing relevant references. In addition, we display the list of the participants with their e-mail addresses.

Acknowledgments

We would like to thank all participants for their effort to travel to Bern and for making “HadAtom05” an exciting meeting. We cordially thank our secretaries, Ruth Bestgen and Ottilia Hänni, for the excellent performance, and the staff of the Institut für Theoretische Physik for their invaluable support. Last but not least, we also thank our colleagues from the program panel, namely Jürg Gasser, Heinrich Leutwyler, Leonid Nemenov, Akaki Rusetsky and Dirk Trautmann, for their continuous help in structuring this workshop.

Bern and CERN, July 2005

Leonid Afanasyev
Gilberto Colangelo
Jürg Schacher

References

- [1] Proceedings of the International Workshop “Hadronic Atoms and Positronium in the Standard Model”, Dubna, 26-31 May 1998, Ed. M.A. Ivanov, A.B. Arbuzov, E. A. Kuraev, V.E. Lyubovitskij, A.G. Rusetsky.
- [2] J. Gasser, A. Rusetsky and J. Schacher, “HadAtom99”, arXiv:hep-ph/9911339.
- [3] J. Gasser, A. Rusetsky and J. Schacher, “HadAtom01”, arXiv:hep-ph/0112293.
- [4] L. Afanasyev, A. Lanaro and J. Schacher, “HadAtom02”, arXiv:hep-ph/0301266.
- [5] J. Gasser, A. Rusetsky and J. Schacher, “HadAtom03”, arXiv:hep-ph/0401204.

2 List of Participants

1	Afanasyev Leonid	Dubna	afanasev@nu.jinr.ru
2	Allkofer Yves	Zurich	Yves.Allkofer@cern.ch
3	Aste Andreas	Basel	aste@quasar.physik.unibas.ch
4	Cabibbo Nicola	Roma	nicola.cabibbo@roma1.infn.it
5	Colangelo Gilberto	Bern	gilberto@itp.unibe.ch
6	dos Santos Covita Daniel	PSI	daniel.covita@psi.ch
7	Dreyer Ute	Basel	dreyer@quasar.physik.unibas.ch
8	Friedman Eli	Jerusalem	elifried@vms.huji.ac.il
9	Gasser Juerg	Bern	gasser@itp.unibe.ch
10	Goldin Daniel	Basel	daniel.goldin@cern.ch
11	Gotta Detlev	Jlich	d.gotta@fz-juelich.de
12	Heim Thomas	Basel	thomas.heim@unibas.ch
13	Hencken Kai	Basel	k.hencken@unibas.ch
14	Ivanov Andrei	Vienna	ivanov@kph.tuwien.ac.at
15	Jensen Thomas	Paris	Thomas.Jensen@spectro.jussieu.fr
16	Johnson Ian	Zurich	Ian.Johnson@cern.ch
17	Juge Jimmy	Dublin	juge@maths.tcd.ie
18	Kubis Bastian	Bonn	kubis@itp.unibe.ch
19	Lemmer Richard	Johannesburg	iclemmer@iafrica.com
20	Lyubovitskij Valery	Tuebingen	valeri.lyubovitskij@uni-tuebingen.de
21	Mannelli Italo	PISA	mannelli@pi.infn.it
22	Marton Johann	Vienna	johann.marton@oeaw.ac.at
23	Minkowski	ITP, Bern	mink@itp.unibe.ch
24	Nemenov Leonid	CERN-JINR	Leonid.Nemenov@cern.ch
25	Pineda Antonio	Barcelona	pineda@ecm.ub.es
26	Raha Udit	Bonn	udit@itkp.uni-bonn.de
27	Rusetsky Akaki	Bonn	rusetsky@itkp.uni-bonn.de
28	Sainio Mikko	Helsinki	mikko.sainio@helsinki.fi
29	Santos, Jos Paulo	Lisboa	jps@cii.fc.ul.pt
30	Sazdjian Hagop	Paris	sazdjian@ipno.in2p3.fr
31	Schacher Juerg	Bern	juerg.schacher@lhcp.unibe.ch
32	Simons Leopold	PSI	leopold.simons@psi.ch
33	Tarasov Alexandr	Dubna	tarasov@nu.jinr.ru
34	Torleif Ericson	CERN	torleif.ericson@cern.ch
35	Trautmann Dirk	Basel	Dirk.Trautmann@unibas.ch
36	Voskresenskaya Olga	Dubna	Olga.Voskresenskaja@mpi-hd.mpg.de
37	Yaz'kov Valeriy	Dubna	Valeri.Iazkov@cern.ch
38	Zemp Peter	Bern	zemp@itp.unibe.ch
39	Zhabitsky Mikhail	Dubna	zhabitsk@nusun.jinr.ru

3 Scientific program

Page

M. Zhabitsky	
<i>The DIRAC experiment at CERN</i>	7
L. Nemenov	
<i>Using $\pi^+\pi^-$ and πK atoms for the experimental check of the low-energy QCD</i>	8
L. Afanasyev and L. Nemenov	
<i>Production of the long-lived excited states of $\pi^+\pi^-$ atoms for DIRAC</i>	9
T. Heim, K. Hencken , M. Schumann, D. Trautmann and G. Baur	
<i>Electromagnetic Exctiation and Ionization of Pionium: An Update</i>	10
J. Juge	
<i>$\pi\pi$ threshold parameters from lattice calculations</i>	12
D. Gotta	
<i>The pionic hydrogen experiment at PSI</i>	13
Ulf-G. Meißner, Udit Raha and Akaki Rusetsky	
<i>The πN scattering lengths from pionic deuterium</i>	14
M.E. Sainio	
<i>Progress report in πN analysis</i>	15
P. Zemp	
<i>Decay width of the ground state of pionic hydrogen in QCD+QED</i>	16
T. E. O. Ericson and A. N. Ivanov	
<i>The π^-p atomic level shift and Isospin Violation in πN Scattering</i>	17
T.S. Jensen	
<i>Kinetic energy distributions in pionic hydrogen</i>	18
A. Pineda	
<i>The chiral structure of the (muonic) hydrogen</i>	19
J. Marton	
<i>Precision Measurements with Kaonic Atoms</i>	20
U. Raha	
<i>Kaonic hydrogen</i>	21
J. Santos	
<i>Kaonic and sigmonic atoms</i>	n.r.
A. N. Ivanov , M. Cargnelli, M. Faber, H. Fuhrmann, V. A. Ivanova, J. Marton, N. I. Troitskaya, and J. Zmeskal	
<i>On the energy level displacement of the ground state of kaonic deuterium</i>	22
Amand Faessler, Th. Gutsche, V. E. Lyubovitskij , K. Pumsa-ard	
<i>Nucleon stucture in Lorentz covariant chiral quark model</i>	23

I. Mannelli	
<i>Study of pion-pion interaction in $K \rightarrow 3\pi$ decays</i>	24
N. Cabibbo	
<i>Cusp effects in $K \rightarrow 3\pi$ and the determination of $\pi\pi$ scattering lengths</i>	n.r.
S. Bakmaev, A. Tarasov , and O. Voskresenskaya	
<i>The unitary correction to the Molière–Fano multiple scattering theory</i>	25
S. Bakmaev and O. Voskresenskaya	
<i>Production of pioniums in the coherent nuclear-nuclear collisions</i>	26
E. Friedman	
<i>Pion-nucleus interaction at low energy</i>	27
J. Gasser	
<i>Concluding remarks</i>	28

The DIRAC experiment at CERN

M. Zhabitsky on behalf of DIRAC collaboration

Joint Institute for Nuclear Research, Joliot-Curie 6, 141980 Dubna, Russia

The aim of the DIRAC experiment at CERN [1] is to measure the lifetime of pionium, a hydrogen-like atom consisting of a π^+ and a π^- meson. The lifetime τ is dominated by the charge exchange process ($\pi^+\pi^- \rightarrow \pi^0\pi^0$) and thus is inversely proportional to the squared difference between S-wave $\pi\pi$ scattering lengths with isospin 0 and 2: $|a_0 - a_2|$. This difference was calculated within the framework of Standard Chiral Perturbation Theory with a precision 1.5% [2] which leads to the lifetime prediction $\tau_{1S} = (2.9 \pm 0.1) \times 10^{-15}\text{s}$.

The DIRAC exploits the lifetime measurement method proposed in [3]. Pionium atoms are produced in proton-nucleus interactions. After production these relativistic atoms may either decay into $\pi^0\pi^0$ or get excited to higher quantum numbers, or break-up in the target material. In the case of break-up, characteristic pion pairs n_A emerge which have a low relative momentum Q in their centre of mass system. The number of produced atoms N_A is proportional to the number of free pion pairs with small Q which undergo Coulomb interaction in final state N_{CC} . Then for a given target material there is a one-to-one correspondence between the measured break-up probability P_{br} and atomic lifetime [4].

In 2001 DIRAC collaboration obtained a first high statistics ($\sim 10^4$) atomic data sample collected from p-Ni interactions at 24 GeV/c [5]. The analysis of experimental distributions on $\pi\pi$ pairs relative momentum results in $n_A^{\text{rec}} = 6530 \pm 294$ pairs from pionium break-up. Then the break-up probability P_{br} becomes [to be published]:

$$P_{\text{br}} = \frac{n_A}{N_A} = \frac{n_A^{\text{rec}}(Q \leq Q_{\text{cut}})}{k(Q_{\text{cut}})N_{CC}^{\text{rec}}(Q \leq Q_{\text{cut}})} = 0.452 \pm 0.023|_{\text{stat}}, \quad (1)$$

which corresponds to the lifetime of the atomic ground state

$$\tau_{1S} = 2.91^{+0.45}_{-0.38}|_{\text{stat}} \, ^{+0.19}_{-0.49}|_{\text{syst}} \times 10^{-15}\text{s}. \quad (2)$$

The systematical error is dominated by uncertainties in multiple scattering, presence of non-identified admixture of K^+K^- and proton-antiproton pairs, uncertainties in the correction due to finite size production and strong interaction for Coulomb-correlated pairs. The analysis of the full data sample (about 2 times more than analysed here) and dedicated measurements, which will decrease systematical errors, is in progress.

References

- [1] B. Adeva et. al., DIRAC proposal, CERN-SPSLC-95-1, SPSLC-P-284 (1995).
- [2] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B603 (2001) 125.
- [3] L. Nemenov, Yad. Fiz. 41 (1985) 980; (Sov. J. Nucl. Phys. 41 (1985) 629).
- [4] L.G. Afanasyev and A.V. Tarasov, Phys. At. Nucl. 59 (1996) 2130; T.A. Heim et al., J. Phys. B33 (2000) 3583; see also references there.
- [5] B. Adeva et al., J. Phys. G30 (2004) 1929.

Using $\pi^+\pi^-$ and πK atoms for the experimental check of the low-energy QCD

L.Nemenov^{1,2}

¹CERN, CH-1211 Geneva 23, Switzerland

²Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

The lifetime (τ) measurement of atoms consisting of $\pi^+\pi^-$ ($A_{2\pi}$) and πK mesons ($A_{\pi K}$) allows to obtain in a model independent way the difference of $\pi\pi$ ($a_0 - a_2$) and πK ($a_{1/2} - a_{3/2}$) scattering lengths in S-state. Basing on the last experimental data on $\tau(A_{2\pi})$ from DIRAC experiment at CERN the expected precisions of this parameter for the approved upgrade of the experiment are presented in Tables 1 and 2. The expected accuracy of $\tau(A_{2\pi})$ and $\tau(A_{\pi K})$ measurement on J-PARC, GSI and SPS CERN are presented in the same tables also. Observation of the long-lived (metastable) states of $A_{2\pi}$ opens the possibility to measure the Lamb-shift in $A_{2\pi}$ and the measurement the other combination the $\pi\pi$ scattering lengths.

Table 1: Statistics of $A_{2\pi}$ (n_A) and the corresponding statistical error (stat) in $|a_0 - a_2|$ to be obtained with the upgraded DIRAC setup during 12 month runs (20h/day) are shown for different accelerators. The latter value should be compared to the contemporary limits in accuracy of $|a_0 - a_2|$ coming from the utilized theoretical dependences of $\tau(A_{2\pi})$ on the $|a_0 - a_2|$ ($\tau = f(a_0 - a_2)$) and the probability of the $A_{2\pi}$ breakup in the target on τ ($P_{br} = f(\tau)$).

	n_A	stat	$\tau = f(a_0 - a_2)$	$P_{br} = f(\tau)$
PS CERN 24 GeV/c	85000	2%	0.6%	1.2%
J-PARC 50 GeV/c	4.1×10^5	0.9%	0.6%	1.2%
GSI 90 GeV/c	1.2×10^6	0.6%	0.6%	1.2%
SPS CERN 24 GeV/c	1.26×10^6	0.5%	0.6%	1.2%

Table 2: The same values as in Table 1 for $A_{\pi K}$ atoms and $|a_{1/2} - a_{3/2}|$ value.

	n_A	stat	$\tau = f(a_{1/2} - a_{3/2})$	$P_{br} = f(\tau)$
PS CERN 24 GeV/c	7000	10%	1.1%	1.2%
J-PARC 50 GeV/c	4.1×10^5	7%	1.1%	1.2%
GSI 90 GeV/c	1.2×10^6	2.5%	1.1%	1.2%
SPS CERN 24 GeV/c	1.26×10^6	2.5%	1.1%	1.2%

Production of the long-lived excited states of $\pi^+\pi^-$ atoms for DIRAC

L. Afanasyev and L. Nemenov

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

Measurement of the $\pi^+\pi^-$ atom lifetime in the DIRAC experiment will allow one to obtain in the model independent way the value $|a_0 - a_2|$, the difference of the s -wave $\pi\pi$ scattering lengths with the isotop spin 0 and 2 correspondently [1]. To get the values of a_0 and a_2 separately basing on the $\pi^+\pi^-$ atom data, one may use the fact that the energy splitting between levels ns and np , $\Delta E_n = E_{ns} - E_{np}$, depends on the another combination of scattering lengths: $2a_0 + a_2$. Thus the measurement the energy splitting coupled with the lifetime measurement would provide a determination of a_0 and a_2 separately [1,2].

The lifetimes of the np states are significantly, 3–5 order, higher in compare with the ground state [2]. For that reason atoms in np states have the mean paths of teens centimeters. Methods of ΔE_n measurement proposed in papers [3] and [4] based on observation of the interference between ns and $np(m=0)$ states in the external electro-magnetic fields.

Production of $\pi^+\pi^-$ atom in the np states have been considered for different target materials and thicknesses with the intent to optimize the experimental conditions for their observation. It have been shown that for the DIRAC experiment usage of thinner targets with smaller Z provides increase of yield of np states and a better ratio to the atom break-up. In the following table a set of targets providing the highest yield of $\pi^+\pi^-$ atom states with the magnetic quantum number $l \geq 1$ are shown.

Target Z	Thickness μm	Break-up	$\sum(l \geq 1)$	$2p_0$	$3p_0$	$4p_0$	$\sum(l = 1, m = 0)$
04	50	2.63%	5.86%	1.05%	0.54%	0.20%	1.93%
06	50	5.00%	6.92%	1.46%	0.51%	0.16%	2.52%
13	20	5.28%	7.84%	1.75%	0.57%	0.18%	2.63%
28	5	9.42%	9.69%	2.40%	0.58%	0.18%	3.29%
78	2	18.8%	10.5%	2.70%	0.55%	0.16%	3.53%

The first estimation of expected accuracy in the energy splitting shows that for the magnetic field of 1 T and 10 cm long measurement during 6 months (3+3) at PS CERN will provide accuracy in about 6% for $\pi^+\pi^-$ atom. For this calculation only atomic pairs was considered as the background.

References

- [1] B. Adeva et al., *Lifetime measurement of $\pi^+\pi^-$ atoms to test low energy QCD predictions*, Proposal to the SPSLC, CERN/SPSLC 95–1, SPSLC/P 284, Geneva, 1995.
- [2] L.L. Nemenov, *Yad.Fiz.* **41** (1985) 980.
- [3] L.L. Nemenov, V.D. Ovsiannikov, *Phys.Lett.* **B 514** (2001) 247.
- [4] L.L. Nemenov, V.D. Ovsiannikov and E.V. Tchapyguine, *Nucl.Phys.* **A 710** (2002) 303.

Electromagnetic Excitation and Ionization of Pionium: An Update

T. Heim¹, **K. Hencken**¹, M. Schumann¹, D. Trautmann¹ and G. Baur²

¹Institut für Physik, University of Basel, Klingelbergstr. 82, 4056 Basel, Switzerland

²Institut für Kernphysik. Forschungszentrum Jülich, 52425 Jülich, Germany

The precise knowledge of excitation and ionization cross sections of pionium going through matter is an important theoretical input into the analysis of the DIRAC experiment [1]. In order to achieve an accuracy of 5% for the lifetime of the pionium the cross section needs to be known to a high precision better than 1%.

At this accuracy a number of additional processes must be taken into account. This has been achieved by our group in a series of papers. The dominant contribution coming from the electric interaction has been studied within a semiclassical treatment in [2]. There, we calculated total and partial cross sections and we studied the influence of atomic screening. The inclusion of target inelastic processes, that is “antiscreeing”, was investigated in [3] and was found to be important (with a correction up to 13%) for light target atoms. The magnetic interaction, the A^2 interaction, and also additional relativistic corrections were found to be below the required 1% level [4]. Multiphoton exchange corrections were calculated within Glauber theory; they reduce the differential cross section by as much as 30% for heavy targets [5,6].

In order to test the accuracy of the Glauber calculations we have tried to calculate some important cross sections within a coupled channel approach [6]. It was found that in a calculation taking only the bound states into account, the effect of higher orders is substantially underestimated. Inclusion of the continuum states by discretizing them with Weyl wave packets was found to give a correction in the right direction, but the overall agreement with the Glauber calculation is still poor. Therefore, as an alternative approach we are now studying the possibility of including the continuum states either within a perturbative treatment, or within a polarisation potential model, where the high lying continuum states can be approximately integrated out. With the help of the closure approximation only matrix elements between bound states need to be evaluated. This work is still in progress.

A peculiarity occurs in calculating the deexcitation cross section, where the pionium undergoes a $2p \rightarrow 1s$ transition while the atom is excited. Due to the relativistic motion one may encounter a singularity of the photon propagator $1/Q^2$. The integration over Q then leads to a cross section that is formally divergent. This problem is not new, of course, going back to Fano [7]. More recently it has been studied in connection with a future muon collider, in the process $\mu^- + \mu^+ \rightarrow e + \bar{\nu} + W^+$ through a ν_μ [8,9] and in connection with atomic physics processes [10]. Different scenarios lead to a regularisation of the photon propagator.

In the case of the DIRAC experiment two phenomena are of importance, having to do either with the finite lifetime of the excited pionium beam, or with the finite lifetime of the propagating photon. The $2p$ state of pionium is not stable but can “decay” either by a transition to the $1s$ state, or by interactions with the target atoms, leading, e.g., to further excited states. Both processes lead to an imaginary part of the mass of the $2p$ state which

then translates to an imaginary part for the photon propagator

$$q^2 \rightarrow q^2 + \frac{i}{2}m\Gamma,$$

with m the mass and Γ the width of the $2p$ states. We find that the width due to the additional interaction is of the order of 1 eV and therefore the dominant effect.

For the photon one gets a finite lifetime due to its attenuation in matter. Starting from the telegraph equation we have derived the modification to the propagator which again acquires an imaginary part, this time of the form

$$q^2 \rightarrow q^2 + \frac{i\Delta}{\lambda c^2},$$

where λ is the attenuation coefficient.

A first study shows that both effects, the pionium lifetime as well as the photon lifetime, are roughly equally important. Whereas these effects can result in changes of the cross sections of the order of a few percent for higher atomic excitation energies, the target inelastic contribution itself constitutes only a small part of the total cross section. Therefore we conclude that the regularisation of the cross section, even though formally needed to get a finite result, does not lead to a change of the cross section at the required accuracy.

References


- [1] B. Adeva *et al.* [DIRAC Collaboration], J. Phys. G **30** (2004) 1929 [arXiv:hep-ex/0409053].
- [2] Z. Halabuka, T. A. Heim, K. Hencken, D. Trautmann and R. D. Viollier, Nucl. Phys. B **554** (1999) 86.
- [3] T. Heim *et al.*, J. Phys. B **33** (2000) 3583
- [4] T. Heim *et al.*, J. Phys. B **34** (2001) 2763
- [5] M. Schumann *et al.*, J. Phys. B **35** (2002) 2683
- [6] M. Schumann, Ph. D. thesis, U Basel (2003) unpublished
- [7] U. Fano, Annu. Rev. Nucl. Sci. **13** (1963) 1
- [8] I. F. Ginzburg, hep-ph/9509314
- [9] K. Melnikov and V. G. Serbo, Phys. Rev. Lett. **76** (1996) 3263, Nucl. Phys. B **483** (1997) 67.
- [10] A. Voitkiv, J. Phys. B **33** (2000) 1299, Physics Reports **392** (2004) 191, and submitted (2005).

Hadronic Scattering in Lattice QCD

K.J. Juge

School of Mathematics, Trinity College, Dublin 2, Ireland

Lattice QCD calculations of the I=2 $\pi\pi$ scattering length was reviewed. Discussions of the various systematic uncertainties involved in lattice calculations ([1]-[5]) were given centered around the most recent $N_f = 2$ simulations of the CP-PACS collaboration [4]. A peculiar extrapolation to the physical pion mass limit (which has also been noted by other collaborations) were discussed using the data from the BGR collaboration who use a chirally symmetric lattice action [2]. The extrapolations are shown in the figures below along with the result from chiral perturbation theory [6].



References

- [1] JLQCD Collaboration, Phys. Rev. **D66** (2002) 077501.
- [2] BGR Collaboration, arXiv:hep-lat/0309075.
- [3] CP-PACS Collaboration, Phys. Rev. **D67** (2003) 014502.
- [4] CP-PACS Collaboration, Phys.Rev. **D70** (2004) 074513.
- [5] BGR Collaboration, arXiv:hep-lat/0409064.
- [6] G. Colangelo *et al.*, Nucl. Phys. **B603** (2001) 125.

The pionic hydrogen experiment at PSI

D. Gotta, for the PIONIC HYDROGEN collaboration

Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany

Experiment R-98.01 [1] performed at PSI aims at a high precision determination of the strong-interaction shift ϵ_{1s} and width Γ_{1s} by means of X-ray spectroscopy of ground-state transitions in pionic hydrogen. From these quantities the πN scattering lengths and the πN coupling constant are extracted [2,3,4,5].

Varying the target density, the observed non-effect on the ground-state transition energy excludes any influence of cascade effects within the experiment's accuracy yielding $\epsilon_{1s} = 7.120 \pm 0.013$ eV [6,7,8]. In order to improve significantly on the accuracy achieved for Γ_{1s} in previous measurements [9] three different X-ray transitions, $\pi H(2p - 1s)$, $\pi H(3p - 1s)$ and $\pi H(4p - 1s)$, have been measured. The degree of Doppler broadening originating from Coulomb de-excitation depends more strongly on the transition than expected. In addition, the $\mu H(3p - 1s)$ transition was measured, where strong-interaction broadening is absent.

The extraction of Γ_{1s} requires a sufficient quantitative understanding of Coulomb de-excitation and an accurate knowledge of the resolution of the crystal spectrometer. The determination of the spectrometer response by using X-rays emitted from an electron-cyclotron resonance ion trap [10] has been addressed by L. Simons [7]. Cascade processes are discussed by T. Jensen [7,11].

From a first analysis of the Doppler broadening in pionic and muonic hydrogen a consistent picture is obtained. In particular, the values extracted for Γ_{1s} for the three pionic transitions agree within the errors. Averaging yields a (preliminary) value of $\Gamma_{1s} = 785 \pm 27$ meV.

References

- [1] PSI experiment R-98.01, <http://pihydrogen.web.psi.ch>.
- [2] V. E. Lyubovitzkij and A. Rusetski, Phys. Lett. B 494 (2000) 9.
- [3] J. Gasser et al., Eur. Phys. J. C 26 (2003) 13.
- [4] P. Zemp, this workshop.
- [5] T. E. O. Ericson et al., Phys. Rev. C 66 (2002) 014005; this workshop.
- [6] M. Hennebach, thesis Universität zu Köln, 2003.
- [7] Proceedings of HadAtom 2003, <http://www.hadatom05.unibe.ch/>.
- [8] D. Gotta et al., Int. J. Mod. Phys. vol 20 (2005) 349.
- [9] H.-Ch. Schröder et al., Eur. Phys. J. C 21 (2001) 433.
- [10] D. F. Anagnostopoulos et al., Nucl. Instr. Meth. B 205, 9 (2003).
- [11] T. Jensen, this workshop.

The πN scattering lengths from pionic deuterium

Ulf-G. Meißner^{1,2}, Udit Raha¹ and Akaki Rusetsky^{1,3}

¹Universität Bonn, Helmholtz-Institut für Strahlen- und Kernphysik (Theorie),
Nußallee 14-16, D-53115 Bonn, Germany

²Forschungszentrum Jülich, Institut für Kernphysik (Theorie),
D-52425 Jülich, Germany

³On leave of absence from: High Energy Physics Institute,
Tbilisi State University, University St. 9, 380086 Tbilisi, Georgia

We use the framework of effective field theories to discuss the determination of the S -wave πN scattering lengths a_+ and a_- from the recent high-precision measurements of pionic deuterium observables [1]. In particular, we investigate the accuracy limits on the calculations of the pion-deuteron scattering length $a_{\pi d}$ in the multiple-scattering series, which are derived in effective field theories. It has been shown that the most stringent limit is set by the presence of the 3-particle low-energy constant (LEC) which encodes the information about the underlying QCD dynamics at a scale $\simeq 1$ GeV. The ways to estimate this constant from fitting to the experimental data, as well as using resonance saturation, have been discussed. We further study in detail the differences between various field-theoretical approaches to the pion-deuteron problem [2,3,4]. After clarifying physical reasons for these (seeming) differences, they have been finally removed from the pion-deuteron data analysis.

The results contained in this talk have been recently published in Ref. [5].

References

- [1] D. Chatellard *et al.*, Nucl. Phys. A **625** (1997) 855; P. Hauser *et al.*, Phys. Rev. C **58** (1998) 1869.
- [2] S. R. Beane, V. Bernard, T.-S. H. Lee and U.-G. Meißner, Phys. Rev. C **57** (1998) 424 [arXiv:nucl-th/9708035]. S. R. Beane, V. Bernard, E. Epelbaum, U.-G. Meißner and D. R. Phillips, Nucl. Phys. A **720** (2003) 399 [arXiv:hep-ph/0206219].
- [3] B. Borasoy and H. W. Griesshammer, Int. J. Mod. Phys. E **12** (2003) 65 [arXiv:nucl-th/0105048].
- [4] S. R. Beane and M. J. Savage, Nucl. Phys. A **717** (2003) 104 [arXiv:nucl-th/0204046].
- [5] U.-G. Meißner, U. Raha and A. Rusetsky, arXiv:nucl-th/0501073.

Progress report in πN analysis

M.E. Sainio

Helsinki Institute of Physics, P.O. Box 64, 00014 University of Helsinki, Finland

There are still quite a few publications appearing in the literature reporting results of pion-nucleon scattering experiments eventhough most of the meson factory experiments have finished their data taking. An exception is the pionic hydrogen experiment which is expected to improve the level shift and width measurements. In 2004 two data analyses have been published, the GWU partial wave analysis [1] and the Bugg analysis [2]. They agree on the value of the pion-nucleon coupling constant producing a number near $f^2 = 0.076$ with an error bar of about 1 %. A slightly higher value, 0.078, is obtained by Ericson et al. [3] in an analysis of the pionic hydrogen data.

The standard for pion-nucleon analysis was set by the work of the Karlsruhe group in the 70's and 80's. In an attempt to update that analysis, we have in Helsinki been working on the expansion method for implementing the fixed- t dispersion relations. As an example for the C^+ amplitude ($C = A + \nu/(1 - t/4m^2)B$) we have

$$C^+(\nu, t) = C_N^+(\nu, t) + H(Z, t) \sum_{n=0}^N c_n^+ Z^n,$$

where H is adjusted to the asymptotic behaviour of the amplitude and

$$Z(\nu^2, t) = \frac{\alpha - \sqrt{\nu_{th}^2 - \nu^2}}{\alpha + \sqrt{\nu_{th}^2 - \nu^2}}.$$

Currently we are still working on data generated with an existing phase shift solution (KA84), but, gradually, the updated data base will be used. In the forward direction the expansion method is simple to use. For the $C^\pm$ amplitudes input for the imaginary parts (total cross sections) and real parts (3 data points for $\text{Re}D^+(\omega, t=0)$) yield an adequate reproduction of the Karlsruhe forward amplitudes, see [4].

References

- [1] R.A. Arndt, W.J. Briscoe, I.I. Strakovsky and R.L. Workman, Phys. Rev. **C69** (2004) 035213.
- [2] D.V. Bugg, Eur. Phys. J. **C33** (2004) 505.
- [3] T.E.O. Ericson, B. Loiseau and S. Wycech, Phys. Lett. **B594** (2004) 76.
- [4] M.E. Sainio, Eur. Phys. J. **A24** (2005), s2, 89.

Decay width of the ground state of pionic hydrogen in QCD + QED

P. Zemp

ITP, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

The width and the shift of the ground state of pionic hydrogen have been calculated in the framework of potential models in ref. [1]. Later the energy shift has been evaluated in a systematic low-energy expansion in [2] whereas the width was considered in the same framework in [3]. Counting $\alpha \sim 1/137$ and $m_d - m_u$ as small quantities of order δ , the result reads

$$\Gamma_{1s} = 8 \alpha^3 M_r^2 p_1^* \left(1 + \frac{1}{P}\right) [a_{0+}^-(1 + \delta_\Gamma)]^2 + \mathcal{O}(\delta^5), \quad (1)$$

with the reduced mass $M_r = M_{\pi^+ m_p} / (M_{\pi^+} + m_p)$, the Panofsky ratio P and the isospin odd scattering length a_{0+}^- . The momentum p_1^* is the center-of-mass momentum of the process $\pi^- p \rightarrow \pi^0 n$. The isospin correction term δ_Γ is

$$\delta_\Gamma = \frac{\epsilon}{a_{0+}^-} + K + \delta_\epsilon^{\text{vac}}, \quad (2)$$

$$K = 4M_r \alpha (1 - \ln \alpha) (a_{0+}^+ + a_{0+}^-) + 2M_r (M_\Sigma - \bar{M}_\Sigma) (a_{0+}^+)^2 + o(\delta). \quad (3)$$

Vacuum polarization effects [4] are contained in $\delta_\epsilon^{\text{vac}}$. The general expression (1) agrees in its form with the one derived by Sigg *et al.* [1] up to a slight difference in the definition of the kinematic prefactor p_1^* . The quantity ϵ can be calculated in CHPT. At tree level calculation [3] yields

$$\delta_\Gamma = (0.6 \pm 0.2) \times 10^{-2} \text{ [CHPT, leading order]}. \quad (4)$$

This result can be compared with $\delta_\Gamma = -(1.3 \pm 0.5) \times 10^{-2}$ found by Sigg *et al.* [1]. A one-loop evaluation of ϵ is underway [5]. For further investigations, see ref. [6].

References

- [1] S. Deser, M. L. Goldberger, K. Baumann and W. Thirring, *Phys. Rev.* **96** (1954) 774; G. Rasche and W. S. Woolcock, *Helv. Phys. Acta.* **49** (1976) 557; D. Sigg *et al.*, *Phys. Rev. Lett.* **75** (1995) 3245.
- [2] V. E. Lyubovitskij and A. Rusetsky, *Phys. Lett. B* **494** (2000) 9 [arXiv:hep-ph/0009206]; J. Gasser, M. A. Ivanov, E. Lipartia, M. Mojzis and A. Rusetsky, *Eur. Phys. J. C* **26** (2002) 13 [arXiv:hep-ph/0206068].
- [3] P. Zemp, "Pionic Hydrogen in QCD + QED: Decay width at NNLO", *PhD Thesis, University of Bern* (2004); J. Gasser and P. Zemp, to be published.
- [4] D. Eiras and J. Soto, *Phys. Lett. B* **491** (2000) 101, [arXiv:hep-ph/0005066].
- [5] P. Büttiker *et al.*, work in progress.
- [6] T. E. O. Ericson, B. Loiseau and S. Wycech, *Phys. Lett. B* **594** (2004) 76; A. N. Ivanov, M. Faber, A. Hirtl, J. Marton and N. I. Troitskaya, *Eur. Phys. J. A* **18** (2003) 653 [arXiv:nucl-th/0306047].

The π^-p atomic level shift and Isospin Violation in πN Scattering

T. E. O. Ericson¹ and A. N. Ivanov²

¹Theory Division, Physics Department, CERN, CH-1211 Geneva 23, Switzerland

² Atomic Institute of the Austrian Universities, Vienna University of Technology, A-1040 Wien and SMI of the Austrian Academy of Sciences, A-1090, Wien

The high precision of data from pionic hydrogen and deuterium permit in principle the determination of the corresponding πN scattering lengths to a precision of up to 0.2% and the e. m. contributions become essential. It is important to have a clear physical picture of their origin and uncertainties. As a complement to the approach via effective field theory with QCD+QED [3] we study the e. m. terms which are generated by the strong interaction, irrespective of its origin. Our starting point is the empirical πN low energy expansion in the absence of the external Coulomb field, which may have intrinsic isospin violating terms. We assume that the e. m. form factors of the pion and nucleon are empirically known and so are the axial form factors. All masses are taken to have the physical values. In short, we have no free parameters. The starting point is the Coulomb problem for the extended charge distribution. The strong interaction at short range is then a perturbation in the spirit of a pseudopotential. By minimal e. m. coupling, or equivalently, interaction energy the πN scattering amplitude has its energy shifted by the Coulomb energy at the origin; this is shown in a model [1]. The wave function at the origin is the one of the extended charge distribution and not the usually used Bohr wave function. In addition a characteristic cusp factor appears to second order in the scattering length, but it is insensitive to assumptions.

All these corrections are of the order of 1% or less with little uncertainty and they are stable with respect to assumptions. The more general case of an extended strong interaction region leads only to weak modifications in practice.

At threshold and in the heavy baryon limit the additional e. m. mechanisms are greatly simplified. Only the dispersive counterpart $\pi N \rightarrow (N, \Delta)\gamma \rightarrow \pi N$ of the Kroll-Ruderman radiative capture process contributes with an axial matrix element and no free parameter [2]. Already the $N\gamma$ intermediate state is an important contribution to the π^-p amplitude, which increases by 9.3% when the intermediate $\Delta\gamma$ states are included degenerate with the nucleon. The isovector amplitude in $m_\pi \ln m_\pi$ agrees with Gasser et al. [3] using only $N\gamma$ intermediate states. This term is nearly canceled by the $\Delta\gamma$ contribution in the limit of the $N\Delta$ mass splitting $\omega_\Delta = 0$. For $\omega_\Delta \neq 0$ the present mechanism contributes 6.5% to the π^-p 1s level shift. These radiative processes are closely governed by the same ΔN physics as in the static limit of πN scattering [4].

References


- [1] T. E. O. Ericson, B. Loiseau, and S. Wycech, Phys. Lett. B **594** (2004) 76.
- [2] T. E. O. Ericson and A. Ivanov, to appear.
- [3] J. Gasser, *et al.*, Eur. Phys. J. C **26**, 13 (2002).
- [4] T. E. O. Ericson and W. Weise, in *Pions and Nuclei*, Clarendon Press, Oxford, 1988.

Kinetic energy distributions in pionic hydrogen


T.S. Jensen

LKB, Ecole Normale Supérieure et Université Pierre et Marie Curie, Case 74,
4 Place Jussieu, F-75252 Paris Cedex 05, France

The goal of the new pionic hydrogen experiment at the Paul-Scherrer-Institut [1] is to measure the line shapes of the $np \rightarrow 1s$ X-ray transitions in π^-p with high precision and to obtain the strong interaction $1s$ shift and width from the results. A difficult problem arises in the case of the extraction of the $1s$ width: the π^-p atoms are in motion at the moment of the radiative transitions so the observed line profiles are broadened due to the Doppler effect. Highly energetic (tens of electron-Volts) π^-p atoms can be produced through Coulomb deexcitation [2]



or by the break-up of excited molecular ion states [3]



Presently, the quantitative understanding of these processes do not allow us to make *ab initio* cascade model predictions [4,5] which are sufficiently accurate for the new experiment. These difficulties have led to the suggestion of an alternative approach for extracting the $1s$ width [6]. The idea consists of combining reliable cascade model input with a fitting procedure which uses parameters to describe the poorly known kinetic energy distributions. Reliable calculations of cross sections are possible for the highly energetic π^-p atoms because the molecular structure of the target hydrogen molecule is less important than for lower energies. This means that the *relative* contributions of for example the $4 \rightarrow 3$ Coulomb component at 73 eV to the broadening of the $3p \rightarrow 1s$ and $2p \rightarrow 1s$ X-ray lines can be predicted. By using these results, a more accurate value of the strong interaction width can be obtained as compared to a fit without constraints.

Acknowledgment: This work was supported by the Swiss National Science Foundation.

References

- [1] D. Gotta, contribution to this workshop.
- [2] L. Bracci and G. Fiorentini, Nuovo Cim. **43A** (1978) 9.
- [3] S. Kilic, J.-P. Karr, and L. Hilico, Phys. Rev. A **70** (2004) 042506.
- [4] T.S. Jensen and V.E. Markushin, Eur. Phys. J. D **21** (2002) 271.
- [5] T.S. Jensen and V.E. Markushin, in *Precision Physics of Simple Atomic Systems*, Eds. S.G. Karshenboim; V.B. Smirnov, Lecture Notes in Physics **627** (Springer, Berlin 2003) 37.
- [6] T.S. Jensen, Eur. Phys. J. D **31** (2004) 11.

The chiral structure of the (muonic) hydrogen

A. Pineda¹

¹Dept. d'Estructura i Constituents de la Matèria, U. Barcelona,
Diagonal 647, E-08028 Barcelona, Catalonia, Spain

High precision measurements in atomic physics provide with a unique place to fix some hadronic parameters related with the proton elastic and inelastic electromagnetic form factors, like the proton radius, magnetic moment, polarization effects, etc.... One complication in this program comes from the fact that widely separated scales are involved in these physical processes. Therefore, it becomes important to relate the physics at these disparate scales in a model independent way. Effective field theories (EFT's) are a natural approach to this problem. In particular, we need an EFT at atomic scales, a natural candidate for which is potential NRQED [1]. By relating this theory with HBET [2], it is possible to obtain the chiral (and model independent) structure of the level splittings of (muonic) hydrogen. Over the last years this idea has been applied in [3]. The matching between HBET coupled to photons and leptons and the relevant EFT at atomic scales have been performed. This matching can be organized in a perturbative expansion in α , $1/m_p$ and the chiral counting. We then computed the $O(m_l^3 \alpha^5 / m_p^2 \times (\ln m_\pi))$ contribution to the Hyperfine splitting (as well as other large logs: $\ln \Delta$, ...) and compared with experiment. This contribution can explain about 2/3 of the difference between experiment and the pure QED prediction when setting the renormalization scale at the ρ mass. The difference can be used to give an estimate of the matching coefficient of the spin-dependent proton-lepton operator in HBET.

The definition of the electromagnetic proton radius has also been studied. It happens to be ambiguous once electromagnetic corrections are considered. This is relevant in view of the possibility to measure the muonic hydrogen Lamb shift at the 30 ppm level at the PSI [4] and, consequently, to obtain the proton radius with an improved accuracy of, at least, one order of magnitude better. We have argued that a natural definition can be given within an EFT framework in terms of a matching coefficient. The definition of the neutron radius was also discussed. The hadronic corrections to the lamb shift (for the polarizability effects only with logarithmic accuracy) were computed within HBET up to $O(m_l^3 \alpha^5 / m_p^2 \times F(m_l / m_\pi))$. It was found that they diverge in the inverse of the pion mass in the chiral limit. Moreover, at this order, they are a prediction of HBET since unknown counterterms are subleading.

References

- [1] A. Pineda and J. Soto, Nucl. Phys. (Proc. Suppl.) **64**, 428 (1998).
- [2] J. Gasser and H. Leutwyler, Annals Phys. **158**, 142 (1984); E. Jenkins and A.V. Manohar, Phys. Lett. **B255**, 558 (1991).
- [3] A. Pineda, Phys. Rev. C **67**, 025201 (2003); hep-ph/0308193; hep-ph/0412142.
- [4] F. Kottmann et al., Hyperfine Interact. **138**, 55 (2001).

Precision Measurements with Kaonic Atoms

J. Marton for the DEAR and SIDDHARTA Collaborations

Stefan Meyer Institut, Austrian Academy of Sciences,
Boltzmanngasse 3, 1090 Wien, Austria

The spectroscopy of X-rays from exotic atoms like pionic and kaonic hydrogen atoms provides precision studies of strong interaction. Kaonic atoms are produced at the DAΦNE electron-positron collider in Frascati. The X-ray transitions in these exotic atoms are studied using an array of CCD X-ray detectors. We obtained the following results:

1) *Kaonic nitrogen*: We measured the X-ray lines and yields of the 7-6, 6-5 and 5-4 transitions for the first time [1]. Kaonic nitrogen is found to be nearly completely stripped of electrons, thus allowing a new kaon mass measurement with a high resolution X-ray detection.

2) *Kaonic hydrogen*: The X-ray spectrum of the kaonic hydrogen K transitions was extracted. The values for the strong interaction shift $\epsilon_{1s} = -193 \pm 37(stat.) \pm 6(syst.)eV$ and width $\Gamma_{1s} = 249 \pm 111(stat.) \pm 30(syst.)eV$ were extracted [2].

The repulsive kaon-proton interaction shown in [3] is verified, but we find smaller values for the shift and width with significantly smaller errors.




Figure 1: Monte Carlo simulated X-ray spectrum of kaonic deuterium using SDDs.

Our experimental program is proceeding toward high precision measurements with new large area X-ray detectors providing timing capability (SDDs) presently in development ¹.

References

- [1] T. Ishiwatari et al., Phys. Lett. B593 (2004) 48.
- [2] G. Beer et al., submitted for publication.
- [3] M. Iwasaki et al., Phys. Rev. Lett. 78 (1997) 3067.

¹EU 6th Framework Program, I3-HP Contract No. RII3-CT-2004-506078

Spectrum and decays of the Kaonic Hydrogen

U.-G. Meißner, **U. Raha**, and A. Rusetsky

*Helmholtz-Institut für Strahlen- und Kernphysik, Universität Bonn
Nußallee 14-16, 53115 Bonn, Germany*

Recent accurate measurements [1,2] of the strong energy shift and the lifetime of the ground state of kaonic hydrogen by DEAR collaboration at LNF-INFN allow one to extract the precise values of the KN scattering lengths from the data. To this end, one needs to relate the latter quantities to the observables of the kaonic hydrogen at the accuracy that matches the experimental precision. In our recent investigations, the problem is considered within the non-relativistic effective Lagrangian approach, which has been previously used to describe the bound $\pi^+\pi^-$, π^-p and πK systems (see, e.g. [3,4,5,6]). We obtain [7] a general expression of the strong shift of the level energy and the decay width in terms of the KN scattering lengths, at $\mathcal{O}(\alpha, m_d - m_u)$ as compared to the leading-order result. It is shown that, due to the presence of the unitarity cusp in the K^-p elastic scattering amplitude above threshold, the isospin-breaking corrections turn out to be very large. This, however, does not affect the accuracy of the extraction of the scattering lengths from the experiment.

References

- [1] R. Baldini et al. [DEAR Collaboration], DAPHNE exotic atom research: The DEAR proposal, LNF-95-055-IR.
- [2] M. Cargnelli et al. [DEAR Collaboration], in Proceedings of the HadAtom03 Workshop, 12-17 October 2003, ECT(Trento, Italy), [arXiv:hep-ph/0401204].
- [3] A. Gall, J. Gasser, V. E. Lyubovitskij, A. Rusetsky, Phys. Lett. B **462** (1999) 335 [arXiv:hep-ph/9905309].
- [4] V. E. Lyubovitskij, A. Rusetsky, Phys. Lett. B **494** (2000) 9 [arXiv:hep-ph/0009206].
- [5] P. Zemp, in Proceedings of HadAtom03 Workshop, 13-17 October 2003 ECT(Trento Italy), J. Gasser, P. Zemp, in preparation, [arXiv:hep-ph/0009206].
- [6] J. Schweizer, Eur. Phys. J. C **36** (2004) 483 [arXiv:hep-ph/0405034].
- [7] U.-G. Meißner, U. Raha, and A. Rusetsky Eur. Phys. J. C **35** (2004) 349 [arXiv:hep-ph/0402261].

On the energy level displacement of the ground state of kaonic deuterium

A. N. Ivanov, M. Cargnelli, M. Faber, H. Fuhrmann, V. A. Ivanova,
J. Marton, N. I. Troitskaya, and J. Zmeskal

Atomic Institute, Vienna University of Technology, Stefan Meyer Institute, Austrian Academy of Sciences, Vienna, Austria and State Polytechnic University of St. Petersburg, St. Petersburg, Russia

The energy level displacement of the ground state of kaonic deuterium is defined by the DGBTT formula¹

$$- \epsilon_{1s} + i \frac{\Gamma_{1s}}{2} = 2\alpha^3 \mu^2 f_0^{K^-d}(0) = 602 f_0^{K^-d}(0), \quad (1)$$

where $\alpha = 1/137.036$ is the fine-structure constant, $\mu = m_K m_d / (m_K + m_d) = 391$ MeV is the reduced mass of the K^-d pair for $m_K = 494$ MeV and $m_d = 1876$ MeV and $f_0^{K^-d}(0)$ is the S-wave amplitude of K^-d scattering at threshold. We show Ref.[1] that the S-wave scattering length $\text{Re } f_0^{K^-d}(0) = a_0^{K^-d}$ is described well by the Ericson–Weise–like scattering length of K^-d scattering Ref.[2]. The imaginary part $\text{Im } f_0^{K^-d}(0)$ we calculate in terms of the contributions of the two-body reactions $K^-d \rightarrow NY$, where $NY = n\Lambda^0$, $n\Sigma^0$ and $p\Sigma^-$, and the experimental data on the two-body reaction rates (see Ref.[33] of Ref. [1]). Our theoretical predictions for the two-body reaction rates agree well with the experimental data. The S-wave amplitude of K^-d scattering at threshold is equal to $f_0^{K^-d}(0) = (-0.540 \pm 0.095) + i(0.521 \pm 0.075)$ fm Ref.[1]. Our prediction for the energy level displacement of the ground state of kaonic deuterium Ref.[1]

$$- \epsilon_{1s} + i \frac{\Gamma_{1s}}{2} = 602 f_0^{K^-d}(0) = (-325 \pm 60) + i(315 \pm 50) \text{ eV} \quad (2)$$

can be used for the planning experiments by the DEAR/SIDDHARTA Collaborations at Frascati (C. Guaraldo *et al.*, *Measurement of kaonic hydrogen with DEAR at DAFNE and future perspectives*, Naples, Italy, November 3–7, 2004 and the report by J. Marton “*Precision measurements with kaonic atoms*” at this Workshop *HadAtom05*).

References

- [1] A. N. Ivanov *et al.*, Eur. Phys. J. A **23**, 79 (2005), nucl-th/0406053.
- [2] T. E. O. Ericson and W. Weise, in *Pions and Nuclei*, Clarendon Press, Oxford, 1988.

¹This is the abbreviation of the Deser–Goldberger–Baumann–Thirring–Trueman formula.

Nucleon structure in Lorentz covariant chiral quark model

Amand Faessler, Th. Gutsche, V. E. Lyubovitskij, K. Pumsa-ard

Institut für Theoretische Physik, Universität Tübingen,
Auf der Morgenstelle 14, D-72076 Tübingen, Germany

We developed a manifestly Lorentz covariant chiral quark model for the study of baryons as bound states of constituent quarks. The approach is based on a non-linear chirally symmetric Lagrangian [1]-[3], which involves effective degrees of freedom - constituent quarks and the chiral (meson) fields. In a first step, this Lagrangian can be used to perform a dressing of the constituent quarks by a cloud of light pseudoscalar mesons and other heavy states using the infrared dimensional regularization of loop diagrams [2]. We calculate the dressed transition operators with a proper chiral expansion which are relevant for the interaction of quarks with external fields in the presence of a virtual meson cloud. These operators are used in the calculation of baryon matrix elements [4].

Our main result is: we perform a model-independent factorization of the effects of hadronization and confinement contained in the matrix elements of the bare quark operators and the effects dictated by chiral symmetry. The calculation of these effects can be done independently. In particular, the computing of matrix elements of bare quark operators can then be relegated to quark models based on specific assumptions about hadronization and confinement. The nucleon form factors have a correct infrared-singular structure dictated by chiral symmetry [3,5]. We reproduce the leading nonanalytic (LNA) contributions to the magnetic moments and the charge and magnetic radii of the nucleons. The LNA contribution to the magnetic moments is proportional to the pseudoscalar meson mass M_P . The nucleon radii are divergent in the chiral limit. The LNA contribution to the charge radii is proportional to the chiral logarithm $\ln(m_N^2/M_P^2)$. The LNA contributions to the magnetic radii are represented by the same logarithm as in the case of the charge radii and by the singular term proportional to $1/M_P$.

This work was supported by the DFG under contracts FA67/25-3 and GRK683. This research is also part of the EU Integrated Infrastructure Initiative Hadronphysics project under contract number RII3-CT-2004-506078.

References

- [1] J. Gasser, M. E. Sainio and A. Svarc, Nucl. Phys. B **307** (1988) 779.
- [2] T. Becher and H. Leutwyler, Eur. Phys. J. C **9** (1999) 643 [arXiv:hep-ph/9901384]; JHEP **0106** (2001) 017 [arXiv:hep-ph/0103263].
- [3] B. Kubis and U. G. Meissner, Nucl. Phys. A **679** (2001) 698 [arXiv:hep-ph/0007056].
- [4] V. E. Lyubovitskij, T. Gutsche and A. Faessler, Phys. Rev. C **64** (2001) 065203 [arXiv:hep-ph/0105043]; A. Faessler, T. Gutsche, V. E. Lyubovitskij, K. Pumsa-ard, Prog. Part. Nucl. Phys. **54** (2005) (in print).
- [5] M. A. B. Beg and A. Zepeda, Phys. Rev. D **6** (1972) 2912.

Study of pion-pion interaction in $K \rightarrow 3\pi$ decays

Italo MANNELLI

Scuola Normale Superiore PISA - Italy

The NA48/2 experiment at the CERN SPS is searching for direct CP violation in K^+ -decays to three pions. The experiment uses simultaneous K^+ and K^- beams of 60 ± 3 GeV/c momentum which overlap in the fiducial region. During 2003 and 2004 about 4 billions fully reconstructed $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ decays and more than 100 millions $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ have been collected. The results from a study of the $\pi^0 \pi^0$ invariant mass distribution M_{00} in a partial sample of 30 millions decays $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ are reported, showing a cusp-like structure centered at $M_{00} = 2M_{\pi^+}$. The experimental apparatus consists mainly of a magnetic spectrometer, with a momentum resolution $\sigma(p)/p = 1.02\%(+)0.044\%$ and a liquid Krypton calorimeter segmented transversally into 13248 2cm $\times$ 2cm cells. The calorimeter is 27 X_0 thick and has an energy resolution for photons $\sigma(E)/E = 0.032/\sqrt{E}(+)0.09/E(+)0.0042$ (E in GeV) and space resolution $\sigma(x) = \sigma(y) = 0.42/\sqrt{E}(+)0.06$ cm thus allowing the accurate detection of multi-photons events. By imposing the π^0 invariant mass to pairs of photons and requiring the compatibility of a common origin for four-photon events grouped in two π^0 pairs and at the same time the Kaon invariant mass including the additional charged pion detected in the magnetic spectrometer, decays of $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ can be fully reconstructed with negligible background. The resolution in the $\pi^0 \pi^0$ invariant mass near $M_{00} = 279$ MeV is typically 0.5 MeV. The distribution of the events as a function of M_{00}^2 shows a cusp like structure around $M_{00}^2 = 4M_{\pi^+}^2$ which is definitely not due to an instrumental effect, given the intrinsic smoothness of the acceptance. It is impossible to fit the distribution with the usual parametrization employed by the PDG whenever the cusp region is included in the range for the fit. A good fit, with $\chi^2 = 149/147$ d.o.f., is obtained by using the expression of the matrix element as specified in [1] which takes into account one and two-loops pion-pion scattering, in particular $\pi^+ \pi^- \rightarrow \pi^0 \pi^0$ charge exchange, and allowing for the contribution of the formation of pionium atoms, with their subsequent $\pi^0 \pi^0$ decay, with a branching ratio as calculated by Silagadze [2]. The main ingredient of the fit is the difference $a_0^0 - a_2^0$ between the I=0 and I=2 S-wave pion-pion scattering lengths. With the present sample the statistical error in the determination of $a_0^0 - a_2^0$ is about 4% while the systematic uncertainty is still under evaluation.

In conclusion the study of the M_{00}^2 distribution in $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ decays by the NA48/2 experiment allows for a novel and potentially more precise determination of $a_0^0 - a_2^0$ which is made possible in particular by the quality of its multi-photon detection capability.

References

- [1] N. Cabibbo and G. Isidori, arXiv:hep-ph/0502130.
- [2] Z. K. Silagadze, JETP Lett. **60** (1994) 689 [arXiv:hep-ph/9411382].

The unitary correction to the Molière–Fano multiple scattering theory

S. Bakmaev, **A. Tarasov**, and O. Voskresenskaya

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

The Molière–Fano multiple scattering theory [1,2] is the most used tool for the accounting of the charged particles' multiple scattering effects at experimental data processing. Estimation of the theory accuracy is of especial importance in the case of DIRAC experiment for it's high angular resolution. One possible source of the inaccuracy of M–F theory is the usage in [1] of the approximate expression for the amplitude of elastic scattering of charged particles by atoms, which violates the optical theorem (unitarity condition). We have estimated the relative corrections the the parameters of M–F theory, resulting from restoring of unitarity in the particle–atom scattering theory and found that they are of order $Z\alpha^2$, where Z is the atomic number of target atoms.

References

- [1] G. Molière, Z.Naturforsch, **3a** (1948) 78
- [2] U.Fano, Phys.Rev. **93**(1954) p.117

Production of pioniums in the coherent nuclear-nuclear collisions

S. Bakmaev and O. Voskresenskaya

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

We consider the production of nP -states of pionium atoms in the coherent collisions of a projectile A_p and target A_t nuclei

$$A_p + A_t \rightarrow A_p + A_t + A_{2\pi}(nP) \quad (1)$$

as the possible source of $A_{2\pi}(nP)$ beam for the pionium Lamb-shift measurements [1].

Since the quantum numbers of pionium in nP -states are the same as photon's ones the coherent photoproduction process

$$\gamma + A_t \rightarrow A_{2\pi}(nP) + A_t \quad (2)$$

is quite intense in GeV-region. Another advantage of the coherent reaction (2) is the sharp angular distribution of produced pioniums.

The source of “effective” photons could be provided by projectile nuclear A_p . For the crude estimation of $A_{2\pi}(nP)$ yields the following formula can be used.

$$\omega \frac{d\sigma}{d\omega} = n_{eff}(Z_p, \omega) \alpha^6 (\sigma_{\pi A_t}^{tot} m_\pi)^2 \left(\frac{1}{n^3} - \frac{1}{n^5} \right) \quad (3)$$

Here n_{eff} is the number of “effective” photons with the energy $\omega = E(A_{2\pi})$ produced by projectile nuclear with charge Z_p [2] and n is the principal quantum number of pionium in the nP -state.

References

- [1] L. Nemenov, talk at HadAtom03, arXiv:hep-ph/0401204
- [2] C. Bertulani, G. Baur, Electromagnetic processes in relativistic heavy ion collisions, Physics Report vol. 163 (1988) 299–408

Pion-nucleus interaction at low energy

E. Friedman ¹

Racah Institute of Physics, The Hebrew University, Jerusalem, Israel

An over-view of the pion-nucleus optical potential, as obtained from global fits to large data sets encompassing the whole of the periodic table is presented. Data bases of well over 100 points have existed since the 1990's, and some 10 points due to 'deeply bound' pionic atom states have been added in recent years. Interest in the pion-nucleus interaction at low energies has been focused on the s -wave part of the pion-nucleus optical potential, where the so-called 'anomalous' repulsion of the s -wave pionic atom potential is the empirical finding that the strength of the repulsive s -wave potential inside nuclei is nearly double the values expected on the basis of the free πN interaction. Large scale fits to pionic atom data showed [1] that the model by Weise [2] where the *isovector* scattering amplitude becomes density dependent, due to a chiral-motivated density dependence of the pion decay constant, is capable of removing the anomaly. In an alternative approach the empirical energy dependence of the *isoscalar* amplitude, within the minimal substitution requirement [3] of $E \rightarrow E - V_c$, where V_c is the Coulomb potential, was also found [4] to remove the anomaly. Both models yield equally good fits to the data.

The study of the pion-nucleus interaction was extended above threshold in recent precision measurements of the elastic scattering of 21 MeV $\pi^\pm$ by several nuclei [5]. The purpose was to see if the anomaly is observed also above threshold. In the scattering scenario, unlike in the atomic case, one can study both charge states of the pion, thus increasing sensitivities to isovector effects and to the energy dependence of the isoscalar amplitude due to the Coulomb interaction. Elastic scattering experiments were performed on targets of Si, Ca, Ni and Zr and reasonably good fits to the data were obtained with the conventional pion-nucleus potential, which again displayed the 'anomalous' repulsion. Greatly improved fits were obtained when the energy dependence was included, but some of the anomaly still remained. With the density dependence mentioned above the anomaly disappears, and including also the energy dependence the fits are very good and the anomaly is absent. Thus from the quality of fits the energy dependence is required by the data, and the anomaly is removed if the chiral-motivated density dependence is included in the model.

References

- [1] E. Friedman, Phys. Lett. B **524** (2002) 87; Nucl. Phys. A **710** (2002) 117.
- [2] W. Weise, Nucl. Phys. A **690** (2001) 98c.
- [3] T.E.O. Ericson and L. Tauscher, Phys. Lett. B **112** (1982) 425.
- [4] E. Friedman and A. Gal, Phys. Lett. B **578** (2004) 85.
- [5] E. Friedman et. al., Phys. Rev. Lett. **93** (2004) 122302.

¹Supported by the Israel Science Foundation grant No. 131/01.

Concluding remarks

J. Gasser

ITP, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

I discussed the present status of theory and experiment in Goldstone boson scattering, and relied for this mainly on the very interesting talks presented at the workshop. My personal judgment of the situation is summarized in the table.

Theory		$\longleftrightarrow$	Experiment	
	status	status		status
		$\pi\pi \rightarrow \pi\pi$		
scattering lengths	$\underline{\underline{\checkmark}}$	$\checkmark$	K_{e4}	$\underline{\underline{\checkmark}}$
		$\underline{\underline{\checkmark}}$	$A_{\pi\pi}$	p
		$\checkmark$	$K_{3\pi}$	p
		$\pi N \rightarrow \pi N$		
scattering lengths	$\underline{\checkmark}$	$\checkmark$	$\pi N \rightarrow \pi N$	$\underline{\checkmark}$
		$\underline{\underline{\checkmark}}$	$A_{\pi p}$	$\underline{\checkmark}$
		$\checkmark$	$A_{\pi d}$	p
		$\pi K \rightarrow \pi K$		
scattering lengths	$\underline{\checkmark}$	$\underline{\underline{\checkmark}}$	$A_{\pi K}$	p
		$Kp \rightarrow Kp$		
scattering lengths	p	$\checkmark$ p	A_{Kp} A_{Kd}	$\checkmark$ p

To illustrate the table, the theoretical status of $\pi\pi$ scattering lengths is very good ($\checkmark$), the (theoretical) relation of the scattering lengths to the cusp in $K_{3\pi}$ decays ($\Longleftrightarrow$) needs improvement ($\checkmark$), and the experimental status of determining scattering lengths from $K_{3\pi}$ decays is qualified as “in progress” (p).

Lattice calculations have the potential to provide in the near future reliable information on the various scattering lengths from first principles.

References

- [1] All the speakers of the workshop have contributed to one or several of the topics mentioned in the table - I do not list them individually.